

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

- c) sampling said analogue input signals to derive the cardiac signal (EKG);
  - d) digitising said EKG signal;
  - e) employing wavelet transform analysis to process said digitised EKG signal;
  - f) deriving the wavelet energy surface of the EKG;
  - g) plotting said energy surface against a location parameter  $b$  and a scale parameter;  
and
  - h) visually displaying said signal in real time.
42. (NEW) The method of Claim 41, wherein said wavelet transform analysis uses a continuous wavelet transform discretised for use in the analysis of digitised signals.
43. (NEW) The method of Claim 41, wherein said scale parameter is characterised by a dilation value  $a$ .
44. (NEW) The method of Claim 41, wherein said scale parameter is characterised by a characteristic wavelet frequency, for example the wavelet band pass frequency value  $f_{bpc}$ .
45. (NEW) The method of Claim 41, wherein the step of visually displaying the signal is characterised by a contour plot.
46. (NEW) The method of Claim 41, wherein the step of visually displaying the signal is characterised by a surface plot.

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

47. (NEW) The method of Claim 41, wherein the step of visually displaying the signal is characterised by one type of energy scalogram from a group comprising 2D and 3D energy scalograms.
48. (NEW) A method of decomposition of waveforms in a cardiac signal comprising the steps of:
- a) connecting electrodes to a patient whose heart is in Ventricular Fibrillation (VF);
  - b) deriving analogue input signals from the electrodes;
  - c) sampling said analogue input signals to derive the cardiac signal (EKG);
  - d) digitising said EKG signal;
  - e) employing wavelet transform analysis to process said digitised EKG signal;
  - f) extracting key features from the wavelet transform representation; and
  - g) guiding a resuscitation protocol, said guidance comprising the steps of;
  - h) using an analytical method to determine the likely outcome of a defibrillation shock; and
  - i) determining whether to provide at least one interim therapeutic intervention from a group comprising defibrillatory shock, CPR and pharmaceutical, before shocking.
49. (NEW) The method of Claim 48 wherein the analytical method is characterised by learning vector quantisation (LVQ) methods, for example Kohonen Networks.

61

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

50. (NEW) The method of Claim 48 where the analytical method is characterised by statistical, stochastic methods, for example Bayesian Methods.
51. (NEW) The method of Claim 48 where the analytical method is characterised by multi-layered neural network methods, for example Radial Basis Neural Networks.
52. (NEW) A method of decomposition of waveforms in a cardiac signal comprising the steps of:
- a) connecting electrodes to a presenting patient with a heart in Ventricular Fibrillation (VF);
  - b) deriving analogue input signals from the electrodes;
  - c) sampling said analogue input signals to derive the cardiac signal (EKG);
  - d) digitising said EKG signal;
  - e) employing wavelet transform analysis to process said digitised EKG signal;
  - f) extracting key features from the wavelet transform representation; and
- using an analytical method for determining the optimal time for shocking.
53. (NEW) The method of Claim 52 where the analytical method is characterised by learning vector quantisation (LVQ) methods, for example Kohonen Networks.
54. (NEW) The method of Claim 52 where the analytical method is characterised by statistical, stochastic methods, for example Bayesian Methods.
55. (NEW) The method of Claim 52 where the analytical method is characterised by multi-layered neural network methods, for example Radial Basis Neural Networks.

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

56. (NEW) A method of decomposition of waveforms in a cardiac signal comprising the steps of:

- a) connecting electrodes to a presenting patient whose heart is in Ventricular Fibrillation (VF) after the commencement of Cardio-Pulmonary Resuscitation (CPR);
- b) deriving analogue input signals from the electrodes;
- c) sampling the analogue input signals to derive the cardiac signal (EKG);  
digitising said EKG signal; and
- d) employing wavelet transform analysis to process said digitised EKG signal.

57. (NEW) The method of claim 56 further including the steps of:

- a) filtering said cardiac signal such that the CPR component is disassociated/separated from the heart signal;
- b) producing an energy wavelet scalogram; and
- c) temporally filtering the scalogram using ridge following techniques.

Q1 58. (NEW) The method of claim 57 where said ridge following techniques are characterised by modulus maxima techniques.

59. (NEW) The method of Claim 57 and further including steps for guiding resuscitation protocol, comprising:

- a) extracting key features from the wavelet transform representation

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

- b) using an analytical method for determining the likely outcome of a defibrillation shock; and
  - c) determining whether to provide at least one interim therapeutic intervention from a group comprising immediate defibrillatory shock and CPR, before shocking.
60. (NEW) The method of Claim 59 where said analytical method is characterised by learning vector quantisation (LVQ) methods, for example Kohonen Networks.
61. (NEW) The method of Claim 59 where said analytical method is characterised by statistical, stochastic methods, for example Bayesian Methods.
62. (NEW) The method of Claim 59 where said analytical method is characterised by multi-layered neural network methods, for example Radial Basis Neural Networks.
63. (NEW) A method of decomposition of waveforms in a cardiac signal comprising the steps of:
- a) connecting electrodes to a presenting patient whose heart is in Atrial Fibrillation (AF);
  - b) deriving analogue signals from said electrodes;
  - c) sampling the analogue input signals to derive the cardiac signal (EKG);
  - d) digitising said EKG signal; and
  - e) employing wavelet transform analysis to process said digitised EKG signal.

Q1

Applicant: Addison, Paul Stanley, et al.  
Serial No. 09/980,770  
Filed: 11/01/2001  
Title: METHOD OF ANALYSIS OF MEDICAL SIGNALS

Examiner: To be assigned  
Group: 3737

64. (NEW) The method of claim 63 further including the step of filtering said cardiac signal such that the QRS complex and T components are disassociated/separated from the heart signal, comprising:
- a) producing an energy wavelet scalogram; and
  - b) temporally filtering the scalogram using ridge following techniques.
65. (NEW) The method of claim 64 where said ridge following techniques are characterised by modulus maxima techniques.
66. (NEW) The method of Claim 64 further including steps for guiding the course of therapeutic intervention taken, comprising:
- a) extracting key features from the wavelet transform representation;
  - b) using an analytical method for determining the likely outcome of a cardioversion shock; and
  - c) determining whether to at least one therapeutic intervention from a group comprising cardioversion shock, and drug therapy.
67. (NEW) The method of Claim 66 where said analytical method is characterised by learning vector quantisation (LVQ) methods, for example Kohonen Networks.
68. (NEW) The method of Claim 66 where said analytical method is characterised by statistical, stochastic methods, for example Bayesian Methods.
69. (NEW) The method of Claim 66 where said analytical method is characterised by multi-layered neural network methods, for example Radial Basis Neural Networks.